

Solving inverse problems by combination of maximum entropy and montecarlo simulation

Jan Naudts

*Departement Natuurkunde, Universiteit Antwerpen, UIA,
Universiteitsplein 1, 2610 Antwerpen, Belgium*

(Dated: November 2000)

Abstract

The montecarlo method, which is quite commonly used to solve maximum entropy problems in statistical physics, can actually be used to solve inverse problems in a much wider context. The probability distribution which maximizes entropy can be calculated analytically by introducing Lagrange parameters. The problem of fixing these lagrangean parameters is circumvented by introduction of a microcanonical ensemble which describes a system together with its heath bath. Some further simplifying assumptions make it feasible to do montecarlo sampling of the probability distribution. The method is applied to the example of determining the distribution of the density of the earth from three data. Advantages of the method are guaranteed convergence and a clear information-theoretic foundation.

Maximum entropy has become one of the dominant methods for solving inverse problems of the underdetermined type. The equilibrium probability distribution function (pdf) p of a canonical ensemble of classical statistical physics is known [1] to be the solution of an inverse problem: It maximizes entropy $S(p)$ given some constraints, e.g. given that the expectation value $\langle H \rangle$ of the energy functional $H(x)$

$$\langle H \rangle = \int_X dx p(x) H(x) \quad (1)$$

has some predetermined value U . Now, montecarlo simulation is a well established numerical method, used in models of statistical physics [4] to sample the equilibrium pdf. The strength of the method follows from Markov chain properties which imply that, under the mild condition of ergodicity of the model, the simulation results converge to the exact result. In addition, because of the equipartition theorem[3, 6], one can expect that the configurations generated during the simulation are *typical* solutions of the inverse problem of finding configurations which satisfy the given constraints. The present paper shows that this combination of maximum entropy and montecarlo simulation, which works so well in statistical physics, can be applied to a more general class of inverse problems.

The maximum entropy method is often used to determine a density function ρ defined over some index set I . A toy example is the prediction of the density of the earth as a function of the distance r to the center of the earth from three data: mass M , radius R and moment of inertia J . Other examples are the restoration of images and computer-assisted tomography (CT). See [5]. In such cases it is common practice to interpret ρ , after suitable normalization, as a pdf which can be determined by the maximum entropy method. This is *not* the approach which is followed here. The alternative requires to consider pdfs on the abstract space X of all possible density distributions ρ . At first sight the latter approach may seem impractical for numerical evaluation because of the huge number of degrees of freedom that can be involved. However, by means of montecarlo simulation it is feasible to sample the pdf so that its average and covariance can be determined numerically without ever having to evaluate actual probabilities. At each moment only one density distribution is stored in the memory of the computer.

The formal solution of the maximum entropy problem can be obtained analytically by introducing Lagrange parameters. A remaining problem is that of determining these lagrangean parameters. They have to be tuned so that constraints of the type $\langle H \rangle = U$ are

satisfied. It is shown below that this problem can be avoided by solving the maximum entropy problem in the microcanonical ensemble instead of in the canonical or grand canonical ensemble.

The main advantage of the new method is that it is based on clean theoretical concepts. The disadvantage is that computation times can be large. Therefore the method is complementary to existing techniques, mostly of iterative nature, which are fast but not optimal. A solution obtained with any of these techniques can be taken as starting point for improvement by montecarlo simulation, e.g. to eliminate artifacts from reconstructed images.

In order to fix notations, consider the general problem of a classical experiment consisting of a number of measurements. The experiment is modeled as a function f from some space X of physical variables into some space V of all possible outcomes of the experiment. Given the experimental outcome v , one can define a pdf p on X indicating with which probability points of X can give rise to the experimental outcome v . This pdf is usually non-unique because the experiment yields only a finite amount of information. At this point the *maximum entropy principle* comes into play: From all pdfs that are compatible with the experimental outcome one should select the one which maximizes entropy.

The entropy of the pdf p is given by

$$S(p) = - \int_X dx p(x) \ln p(x) \quad (2)$$

It has to be maximized under the constraints that the averages $\langle f_j \rangle$ equal the experimental data v_j , for $j = 1, 2, \dots, N$. It is straightforward to show that this leads to the result

$$p(x) = \frac{e^{-\sum_j \gamma_j f_j(x)}}{\int_X dx e^{-\sum_j \gamma_j f_j(x)}} \quad (3)$$

with lagrangean parameters γ_j , one for each component f_j of f . These γ_j have to be chosen such that the constraints $\langle f_j \rangle = v_j$ hold. The latter can be a hard problem. It is avoided below by reformulating the problem in a different ensemble.

The description of a mechanical system in canonical or grand canonical ensemble is known to be equivalent to the description of a system interacting with its environment. Let us therefore postulate that a pdf $\rho(x, \omega)$ exists which combines the state x of the system and the noise ω of the environment. Both p and the pdf μ of the environment can be derived

from ρ by

$$p(x) = \int_{\Omega} d\omega \rho(x, \omega) \quad (4)$$

and

$$\mu(\omega) = \int_X dx \rho(x, \omega) \quad (5)$$

The outcome $f(x, \omega)$ of the experiment depends now on both the state of the system and the noise of the environment. The maximum entropy principle of the microcanonical ensemble, containing both system and environment, states that p should be varied in such a way that the entropy is maximal under the constraints that (5) is satisfied and that $f(x, \omega) = v$.

Let us now assume that $f(x, \omega) = v$ has a unique solution as an equation in ω . Denote it $\omega(x, v)$. Then the constraint $f(x, \omega) = v$ implies that

$$\rho(x, \omega) = p(x) \delta_{\omega(x, v)}(\omega) \quad (6)$$

(δ_a is the distribution concentrating in the point a). A straightforward calculation gives

$$p(x) = \frac{1}{Z} e^{-\beta(\omega(x, v))} \quad (7)$$

with Z a normalization factor, and with lagrangean parameters $\beta(\omega)$, one for each possible value of the noise ω . The parameters $\beta(\omega)$ have to be chosen in such a way that (5) holds. This can be done easily. Note that (4) is fulfilled by construction. Using (6, 7) equation (5) becomes

$$\mu(\omega) = e^{-\beta(\omega)} \frac{1}{Z} \sigma_v(\omega) \quad (8)$$

with

$$\sigma_v(\omega) = \int_X dx \delta_{\omega(x, v)}(\omega) \quad (9)$$

There follows

$$p(x) = \frac{\mu(\omega(x, v))}{\sigma_v(\omega(x, v))} \quad (10)$$

In this result the pdf μ is unknown, but is assumed to be fixed by the experimental environment.

Assume now that i) the noise space Ω coincides with the space V of outcomes, ii) the function f is of the form

$$f(x, \omega) = g(x) + \omega \quad (11)$$

iii) μ is of the form

$$\mu(\omega) = (2\pi)^{-k/2} \prod_{j=1}^k \sigma_j^{-1} e^{-(1/2)\omega_j^2/\sigma_j^2} \quad (12)$$

with parameters $\sigma_1 \cdots \sigma_k$ controlling the amount of noise. Let

$$\phi_g(w) = \int_X dy \prod_j \delta(g_j(y) - w_j) \quad (13)$$

with δ Dirac's delta-function. Then one has $\sigma_v(\omega(x, v)) = \phi_g(g(x))$. Using $\omega(x, v) = v - g(x)$ one obtains from (10) our main result

$$p(x) = \frac{1}{(2\pi)^{k/2} \prod_{j=1}^k \sigma_j} \exp(-H(x) - \log \phi_g(g(x))) \quad (14)$$

with

$$H(x) = \frac{1}{2} \sum_{j=1}^k (v_j - g_j(x))^2 / \sigma_j^2 \quad (15)$$

The quantity $H(x)$ is the analog of the energy of statistical mechanics, the quantity $\log \phi_g(w)$ is the entropy, in the sense of Boltzmann, of the macrostate consisting of all physical states y for which $g(y) = w$.

The measurement functions $g_1, \dots, g_k$ can be completed with functions $g_{k+1}, \dots, g_N$ in such a way that together the functions $g_j, j = 1 \cdots N$ form a new coordinate frame for the space X . It is now straightforward to calculate moments of the measurement functions

$$\begin{aligned} \langle g_j^m \rangle &= \int_X dx p(x) g_j^m(x) \\ &= \int dg_j g_j^m (2\pi\sigma_j)^{-1/2} e^{-(1/2)(v_j - g_j)^2/\sigma_j^2} \end{aligned} \quad (16)$$

This means that the pdf (14) is such that each of the variables $g_j, j = 1 \cdots k$, has a normal distribution with average v_j and spread σ_j .

Let us discuss in what follows how (14) can be sampled using montecarlo simulation. Due to the quadratic nature of the hamiltonian H the only contributions to (14) come from points x such that $g(x)$ is close to v (let us assume to this point that the uncertainties σ_j are small so that H becomes large if $g(x)$ is not close to v). Hence, to zeroth order the term $-\log \phi_g(g(x))$ in (14) is constant and can be neglected. For each update $x \rightarrow x'$, considered during execution of the montecarlo algorithm, one has to evaluate the change in energy

$$\begin{aligned} \Delta H &\equiv H(x') - H(x) \\ &= \frac{1}{2} \sum_j (g_j(x') - g_j(x)) [g_j(x') + g_j(x) - 2v_j] \end{aligned} \quad (17)$$

The update is accepted if a random number r , uniformly distributed in $[0, 1]$, is smaller than $\exp(-\Delta H)$. The algorithm makes sense when the calculation of $g(x') - g(x)$ can be done efficiently because then also ΔH can be obtained efficiently. An estimate of the relative vector $g(x) - v$ can be maintained during the simulation by adding $g(x') - g(x)$ to $g(x) - v$ whenever the update is accepted. The estimate can be kept accurate by explicitly evaluating $g(x) - v$ at regular time intervals.

The main effect of the zeroth order approximation, assuming $-\log \phi_g(g(x))$ to be constant in (14), is a violation of (16). In particular the average of $g(x)$, when calculated with the pdf obtained by the montecarlo simulation, will not coincide with v . Let us show how one can correct this deficiency. An expansion to first order gives

$$\log \phi_g(g(x)) \simeq \log \phi_g(v) + a_g(v) \cdot (g(x) - v) + \dots \quad (18)$$

with $a_g(v) = \phi_g(v)^{-1} \nabla \phi_g(v)$. Introduce a new hamiltonian H' by

$$\begin{aligned} H'(x) &\equiv H(x) + \log \phi_g(g(x)) \\ &\simeq \frac{1}{2} \sum_{j=1}^k (v'_j - g_j(x))^2 / \sigma_j^2 + \text{constant terms} \end{aligned} \quad (19)$$

with $v'_j = v_j - (a_g(v))_j \sigma_j$. It is easy to estimate $v' - v$ numerically. Indeed, from the pdf obtained by montecarlo simulation one can calculate the average $\langle g(x) \rangle$ and the deviation w from the target v , i.e.

$$w = \langle g(x) \rangle - v \quad (20)$$

Then a good guess is $v' = w - v$. The montecarlo simulation can now be continued with v replaced by v' . In this way the error can be reduced to second order.

Let us shortly discuss the example of calculating the density of the earth as a function of the distance r to the center of the earth from three data: mass M , radius R and moment of inertia J . The experimental data are $3M/4\pi R^3 = 5517 \pm 5 \text{ kg/m}^3$ and $15J/4\pi R^5 = 0.84 \times 5517 = 4634.28 \pm 55 \text{ kg/m}^3$ – the error bars are simple guesses made by the author. The sphere with radius R is divided into N shells of equal volume. N was chosen equal to 100 as in [5]. The average density of the n -th shell is denoted ρ_n . It satisfies

$$\begin{aligned} \sum_{n=1}^N \rho_n &= N \frac{3M}{4\pi R^3} \\ \sum_{n=1}^N \rho_n (n^{5/3} - (n-1)^{5/3}) &= N^{5/3} \frac{15J}{4\pi R^5} \end{aligned} \quad (21)$$




FIG. 1: Density ρ_n as a function of shell number n

The simulation is started with a uniform mass distribution. Two types of updates are used, an exchange of density of two randomly chosen shells, and an increment/decrement of the density of a randomly selected shell. The two update technique occur with equal probability. The increment/decrement is randomly selected from an interval with self-adapting boundaries: the interval grows by a factor of 1.1 on success, and shrinks by a factor of 0.95 on failure. The simulation times have been chosen excessively large to stress the advantage of the present method that the convergence can only improve. First, 10,000 montecarlo steps (mcs) are used to be sure that the configuration of densities is typical (as usual one mcs contains N update trials). The next 30,000 mcs are used to calculate the averages $\langle g(x) \rangle$ and the deviation w given by (20). The result is

$$\langle g_1(x) \rangle = 5517.19, \quad \langle g_2(x) \rangle = 4673.19 \quad (22)$$

Then the target v is replaced by v' and the simulation runs for 100,000 mcs to determine the average densities $\langle \rho_n \rangle$ – see fig. 1. Due to the correction from v to v' the relations (16)

are better satisfied. Indeed, one obtains

$$\langle g_1(x) \rangle = 5516.97, \quad \langle g_2(x) \rangle = 4635.59 \quad (23)$$

The predicted density in the centre of the earth is about 12,460 kg/m³, at the surface about 3,400 kg/m³. These values should be compared with the results of [5]: 11,200 or 13,600 kg/m³ in the centre of the earth, depending on the method being used, and 3,250 kg/m³ at the surface.

One concludes from the example that the new method works. It can compete with existing numerical techniques on two grounds: i) Absolute convergence. Most real world inverse problems are ergodic, in which case the montecarlo simulation can only improve with increasing computing time. In most iterative methods results start to deteriorate when the computations are not stopped in time. ii) Clear interpretation of the results. What one calculates is the average and variation of that distribution of density functions which maximizes entropy and hence contains the least information. From an information-theoretic point of view this is the best one can do. Of course, any method has its limitations. They will show up when the method is tried to a variety of problems. In particular, one can expect that in problems of image reconstruction montecarlo simulation will be useful to improve solutions obtained by iterative methods. Much can be learned from the experience with montecarlo simulations acquired in statistical mechanics. In particular, it is obvious that efficient techniques for updating configurations are essential for improving computational speed.

-
- [1] E.T. Jaynes, *Information theory and statistical mechanics*. Part I. Phys. Rev. **106**, 620-630 (1957); Part II. Phys. Rev. **108**, 171-191 (1957).
 - [2] E. Rietsch, *The Maximum Entropy Approach to Inverse Problems*, J. Geophys. **42**, 489-506 (1977).
 - [3] R.M. Gray, *Entropy and information theory* (Springer Verlag, 1990)
 - [4] K. Binder and D.W. Heermann, *Montecarlo simulation in statistical physics* (Springer Verlag, 1988)
 - [5] R.M. Bevensee, *Maximum entropy solutions to scientific problems* (Prentice Hall, 1993)

- [6] J.T.Lewis and C.-E. Pfister, *Thermodynamic probability theory: some aspects of large deviations*, Uspekhi Mat. Nauk 50:2 47-88 (1995); Russian Mathematical Surveys 50:2 279-317 (1995).